

A Comprehensive Review of the Impact of Thermal Radiations on Energy Exchange Systems

Muhammad Ashraf

Department of Mathematics, Faculty of Science, University of Sargodha, Sargodha, Pakistan

ABSTRACT

Radiations or electromagnetic rays have a significant role in energy emission from nuclear explosions as well as energy transmission in furnaces and combustion chambers. The radiations are considered in computing thermal influence in devices such as a rocket nozzle, nuclear power plants for space applications, or a gaseous-core nuclear rocket. Energy is transferred between separated elements without a medium in case of radiations or electromagnetic rays. The structure, surface quality, temperature and wavelength of the rotation angle at which radiation is emitted or absorbed by the surface and the spectral distribution of the radiation encountered on the surface are only a few of the variables that affect the radiative behavior. In some important situation, heat is transferred within the solid media or highly viscous media is considered so that convection in the medium is not important. The combined radiation-conduction is important in glassy materials and can be used as a coating to protect the interior of a body from high external temperature. Also, the radiation-conduction process plays a crucial role in controlling how hot or cold the ablating layer is. In these circumstances, the temperature distribution has an impact on how the ablating substance will melt, soften or evaporate. The main novelty of the current work is to highlight the important aspects of thermal radiation effects on heat transfer mechanism and to focus on recent developments in this field.

Keywords: Thermal Radiation, Radiation-conduction Parameter, Radiative Heat Flux, Temperature Field

Classical and Recent Efforts

Present study comprises on two parts that is (i) classical and (ii) recent knowledge of the different aspects of the radiation-conduction heat transfer processes. Special emphasizes is given to the proper usage and determination of radiation-conduction parameter effects on the behavior of heat transfer mechanism which exists in literature. Furthermore, the connection between radiation and the other heat transport modes is discussed with the reference of previous published classical and recent literature. In order to obtain solutions for radiation-conduction energy in an emitting and absorbed medium, the energy equation for two dimensional boundary layer flows by following [6] is of the form:

$$\frac{\partial T}{\partial t} = \nabla \cdot (k \nabla T - q_r) \quad (1)$$

where q_r is the radiative heat flux and can be used to describe the space between two dark panels [6].

$$q_r = - \frac{4\sigma}{3\alpha_R} \frac{\partial T^4}{\partial y} \quad (2)$$

There are two zones that make up the temperature field. As the area typically has a low thickness, it may be assumed that it is optically thin and that radiation will travel through it without being attenuated. The reader interested in this topic is referred to study [6] for further information. Our main focus on current study is to review the influence of radiation-conduction parameter on thermal boundary layer and heat transfer processes. The detail of literature review targeted the above issue is given in below paragraphs.

The injection of absorbing particles is useful to shield the surface of thermal radiation beam if the injection rate is relatively high [1]. In addition to the radiative heat index $Rd \gg 1$, the heat flux in the inner zone of a thick boundary layer cannot be accurately anticipated using the thicker

assumption alone. In comparison to the case where there is no radiation, the presence of radiation tends to diminish enthalpy rates, their inclinations and a thickening of the heat flow separation [3]. The choice of appropriate range of the parameters involved in the flow model and due to the effect of the absorbing radiation at the entrance of the pipe increases the heat transfer rate [4]. For controlled tests of transmitting heat radiation in absorbing and emitting substances, glass is a valuable medium [5]. Because radiant emission is temperature-dependent and thermal radiation is crucial for applications involving the rate of heat and mass transfer [6]. Radiation cannot be disregarded in the prediction of heat exchange because its features rely on the spectral absorption coefficient, index of refraction, thickness, boundary conditions and temperature field [7-8]. In reference [9], the author highlighted the usefulness of a formulation of the radiant energy exchange equation in terms of dimensionless variables for the case in which heat generation rates of the cylinder are known and the temperature are sought. According to Gilpin et al. [10], melting occurs on lakes and streams as a result of assimilation of radiant energy in ice. Since gas absorbs energy, the radiation intensity in wavelength intervals in actual medium drops across the beam path [11].

In each of the three convective processes, the dimensionless local shear stress and local rate of heat transmission are decreased as the thermal radiation parameter is increased [12]. It has been shown that [13], the whole heat transmission in the packed bed under the stated operational circumstances is found to be mostly unaffected by thermal radiation and intra particle conduction. The radiative dispersion hypothesis for radiation heat transfer is the most preferred method [14]. In the presence of thermal radiation, local friction factor is decreased and local Nusselt quantity is increased [15].

*Corresponding author: muhammad.ashraf@uos.edu.pk




Fig. 1: Highlights the velocity profile, temperature distribution and magnetic flux along permeable surface for different values R_d in the presence of aligned magnetic field.

Molla et al. [16] claimed that local Nusselt quantity and the average heat transmission rate is improved when the quantities of radiation/conduction parameter and surface heating parameter along vertical wavy surface are increased. The local velocity, temperature field and magnetic field transversal component all decline when the radiation-conduction index rises [17]. Furthermore, they concluded that for increasing values of radiation parameter the Nusselt number is increased and the obtained results with the help of finite difference method are given in Table 1.

Table 1: Computational data for local Nu_x along the surface distance ξ for two values of radiation parameter R_d by keeping other parameters constant in the flow model under the influence of aligned magnetic field.

ξ	$R_d = 1.0$	$R_d = 10.0$
0.05	0.29162	0.34621
0.1	0.29321	0.34935
0.4	0.30251	0.36789
0.8	0.31429	0.39189
1.0	0.31992	0.40360
4.0	0.39612	0.57401
8.0	0.50613	0.83791
10.0	0.56757	0.98672

Molla et al [18] showed that the shearing force and rate of heat transport are provided as a function of coefficient of skin friction, local mean and Nusselt factor for a wide range of radiation/conduction characteristics. In the presence of thermal radiation, the detail of velocity profile, temperature distribution and current density along a magnetized porous surface has been demonstrated [19]. The obtained numerical

results are presented in Fig. 1(a)-(c). It is shown that, along a porous surface the velocity profile, temperature distribution and magnetic flux are decreased for increasing values of radiation parameter R_d .

The optical and thermal properties of real ceramic foam at high temperature were taken into account both theoretically and experimentally [20].

Radiant heat flux can account for more than one-third of the total thermal gradient in highly porous transparent plastic or polyethylene microspheres. By carefully balancing porosity, one can drastically lower the equivalent conductivity of polymeric foam [21]. The increase of thermal radiation restrains the convective heat transfer and overall heat transfer uniformly [22]. In systems with significant thermal loads, the water jacket acts as a heat shield between hot and cool regions [23]. The non-stationary temperature regime scenario with quick heating and cooling stages, the thickness temperature distribution cannot be predicted by the ETC model with absolute error less than 502013100k [24]. The behavior of thermal radiation in coiled curved pipe for the different ranges of involved parameter is predicted in our previous work [25]. The appearance of metal in functionally graded thermal barrier coating; degraded the insulation performance [26]. In order to control heat radiation, the idea of photon chemical potential is crucial [27]. For increasing values of radiation parameter u and θ are enhanced and ϕ shows the opposite behavior [28]. They obtained the numerical results around different positions of the sphere as given in Fig. 2(a)-(c).




Fig. 2: Highlights the velocity profile(u), temperature distribution (θ) and mass concentration (ϕ) against normal distance Y around the solid sphere for various values of R_d in the presence of aligned magnetic field.

Table 2: Highlight the slopes of dimensionless velocity (u), dimensionless temperature (θ) and mass concentration (ϕ) around different locations of a solid sphere in the presence of aligned magnetic field.

X	$\frac{\partial u}{\partial y}$		$\frac{\partial \theta}{\partial y}$		$\frac{\partial \phi}{\partial y}$	
	$R_d=0.1$	$R_d=1.1$	$R_d=0.1$	$R_d=1.1$	$R_d=0.1$	$R_d=1.1$
$\frac{\pi}{6}$	2.03983	2.12816	0.94670	0.57034	0.37895	0.44799
$\frac{\pi}{4}$	2.47647	2.61447	0.70285	0.42245	0.52161	0.59891
$\frac{\pi}{2}$	2.94238	3.13261	0.53525	0.31783	0.64543	0.72913
π	0.57847	1.26779	-0.41769	1.25722	-1.40916	0.02432

Further, they also concluded the slopes of velocity, temperature and mass concentration for different values of radiation parameter as given in Table 2. It is also shown that the behavior of above said quantities is different at different position.

In associated conductive/radiative heat transfer, the impact of radiant heat grows significantly and contributes for >75% of the heat transmission [29]. The understanding of radiative distribution function characteristics gives a model for the commercial assessment of radiative heat transport in a specific dense system [30].

Conclusion

In current review, we have offered a unified perspective on classical and recent trends towards physical significances of the heat transport consequences of radiant heat on energy exchange systems.

We have summarized the following conclusions:

- All the three modes of the convective heat transfer experience a reduction in locally thermal transport rate and local shear force in the presence of thermal radiation.
- The radiant heat and intraparticle conduction do not play a significant role in the packed bed under specified operating conditions.
- In case of thermal boundary layer, the enthalpy gets lower when radiant energy is contained in the layers.
- Radiative distributive function characteristics provide a reference for industrial analysis and radiative heat transfer in dense particular system.
- The inclusion of thermal radiation or radiative heat flux in diffusion energy equation yields that heat transfer average and local Nusselt number is increased for the increasing values of radiative parameter and surface heating parameter.
- In systems with significant thermal loads, the water jacket acts as a heat shield between hot and cool regions.
- Thus, this review will be useful for the researchers to understand how the classical ideas can tune the current issues raised in energy exchange system?

References

- [1] C.A. Fritsch, R.J. Grosh and M.W. Wildin, "Radiative heat transfer through an absorbing boundary layer," J. Heat Transfer, Vol. 88, no. 3, pp. 296-304, 1966.
- [2] J.L. Novotny and K.T. Yang, "The interaction of thermal radiation in optically thick boundary layers," J. Heat Transfer, Vol. 89, no. 4, pp. 309-312, 1967.
- [3] J. Shawrtz, "Radiation coupled viscous flows," Int. J. Heat and Mass Transfer, Vol. 11, no. 4, pp. 689-697, 1968.
- [4] B.L. Pearce and A.F. Emery, "Heat transfer by thermal radiation and laminar forced convection to an absorbing fluid in the entry region of a pipe," J. Heat Transfer, Vol. 92, no. 2, pp. 221-230, 1970.
- [5] N.D. Eryou and L.R. Glicksman, "An experimental and analytical study of radiative and conductive heat transfer in molten glass," J. Heat Transfer, Vol. 94, no. 2, pp. 224-230, 1972.
- [6] R. Siegel and J.R. Howell, "Thermal radiation heat transfer," NY: McGraw-Hill, 1972.
- [7] E.E. Anderson, R. Viskanata and W.H. Stevenson, "Heat transfer through semitransparent solids," J. Heat Transfer, Vol. 95, no. 2, pp. 179-186, 1973.
- [8] R. Viskanata and E.E. Anderson, "Heat transfer in semitransparent solids, in Thomas F. Irvine, Jr., and James P. Hartnett (eds.)," Advances in Heat Transfer, Vol. 11, pp. 317-441, 1975.
- [9] R.L. Cox, "Radiative heat transfer in arrays of parallel cylinders," Ph.D thesis Tennessee University, Knoxville, 1976.
- [10] R.R. Gilpin, R.B. Robertson and B. Singh, "Radiative heating in ice," J. Heat Transfer, Vol. 99, pp. 227-232, 1977.
- [11] H.A. Vercammen and G.F. Forment, "An improved zone method using Monte Carlo Techniques for the simulation radiation in industrial furnaces," Int. Journal of Heat and Mass Transfer, Vol. 23, no. 3, pp. 329-337, 1980.
- [12] M.A. Hossain and H.S. Takhar, "Radiation effects on mixed convection along a vertical plate with uniform surface temperature," Heat and Mass Transfer, Vol. 31, pp. 243-248, 1996.
- [13] A.A. Jalalzadeh-Azar, W.G. Steele and G.A. Adebijyi, "Heat Transfer in a High-Temperature Packed Bed Thermal Energy Storage System-Roles of radiation and intraparticle conduction," Journal of Energy Resource Technology, Vol. 118, no. 1, pp. 5057, 1996.
- [14] Vadim A. Petrov, "Combined radiation and conduction heat transfer in high temperature fiber thermal radiation," International Journal of Heat and Mass Transfer, Vol. 40, no. 9, pp. 2241-2247, 1997.
- [15] K.A. Yih, "Effect of radiation on natural convection about truncated cone," Int. J. Thermal Science, Vol. 42, no. 23, pp. 4299-4305, 1999.
- [16] M.M. Molla and M.A. Hossain, "Radiation effect on mixed convection laminar flow along a vertical wavy surface," Int. J. of Thermal Science, Vol. 46, no. 9, pp. 926-935, 2007.
- [17] M. Ashraf, S. Asghar and M.A. Hossain, Thermal radiation effects on hydromantic mixed convection flow along a magnetized vertical porous plate. Mathematical Problems in Engineering, vol. 2010, article ID 686596, 30 pages, 2010.
- [18] M.M. Molla, M.A. Hossain and S. Siddiqua, "Radiation effect on free convection laminar flow from an isothermal sphere," Chemical Engineering Communications, Vol. 198, no. 12, pp. 1483-1496, 2011.

- [19] M. Ashraf, S. Asghar and M.A. Hossain, Computational study of the combined effects of conduction radiation and hydro magnetic on natural convection flow past a magnetized permeable plate. *Appl. Math. Mech. (English Edition)*, 33(6), 731-750, 2012.
- [20] R. Coquard, D. Rochais and D. Baillis, "Conductive and radiative heat transfer in ceramic and metal foams at finite temperature," *Fire Technology*, Vol. 48, pp. 699-732, 2012.
- [21] P. Ferlk, R. Pokorney and J. Kosek, "Multiphase approach to couple conduction-radiation heat transfer in reconstructed polymeric foams," *International Journal of Thermal Sciences*, Vol. 83, pp.68-79, 2014.
- [22] G. Yang and J.Y. Wu, "Effects of natural convection, wall thermal conduction, and thermal radiation on heat transfer uniformity at a heated plate located at the bottom of a three-dimensional rectangular enclosure," *Numerical Heat Transfer, Part A: Applications*, Vol. 69, no. 6, pp. 589-606, 2016.
- [23] R. Kantor, "Modeling of a coupled radiation-conduction heat transfer through a heat shield in vacuum thermal isolation applications," *Procedia Engineering*, Vol. 157, pp. 271-278, 2016.
- [24] R.A. Mironov, M.O. Zabezhailov, V.V. Cherepanov, M.Yu. Rusin, "Transient radiative-conductive heat transfer modeling in constructional semitransparent silica ceramics," *International Journal of Heat and Mass Transfer*, Vol. 127, pp. 1230-1238, 2018.
- [25] M. Ashraf, R. Yasmeen and M. Ahmad "Thermal radiation mixed convection boundary layer flow in tightly coiled curved pipe for large value of Richardson number," *Thermal Science*: Vol. 22, No1A, pp. 147-156, 2018.
- [26] W.A. Ge, C.Y. Zhao and B.X. Wang, "Thermal radiation and conduction in functionally graded thermal barrier coatings. Part II: Experimental thermal conductivities and heat transfer modeling," *International Journal of Heat and Mass Transfer*, Vol. 134, pp. 166-174, 2019.
- [27] Li, Y., Li, W., Han, T. *et al.*, "Transforming heat transfer with thermal metamaterials and devices". *Nat Rev Mater* Vol. 6, pp. 488-507, 2021.
- [28] A. Abbas, M. Ashraf and A. Jawad "Chamkha, Combined effects of thermal radiation and thermophoretic motion on mixed convection boundary layer flow," *Journal of Alexandria Engineering*, Vol. 60, pp. 3243-3252, 2021.
- [29] Chao Fan, Xin-Lin Xia, Wei Du, Chuang Sun, Yang Li, "Numerical investigations of the coupled conductive-radiative heat transfer in alumina ceramics," *Heat and Mass Transfer*, Vol. 135, 106097, 2022.
- [30] B. Liu, J. Zhao, L. Liu and A.V. Gusarov, "Analytical relation of radiation distribution function in random particulate systems," *Heat and Mass Transfer*, Vol. 141, 106555, 2023.